

University of Dundee

Precision Medicine in Type 2 Diabetes

Dennis, John M.; Shields, Beverley M.; Hill, Anita V.; Knight, Bridget A.; McDonald, Timothy J.; Rodgers, Lauren R.

Published in:
Diabetes Care

DOI:
[10.2337/dc17-1827](https://doi.org/10.2337/dc17-1827)

Publication date:
2018

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Dennis, J. M., Shields, B. M., Hill, A. V., Knight, B. A., McDonald, T. J., Rodgers, L. R., Weedon, M. N., Henley, W. E., Sattar, N., Holman, R. R., Pearson, E. R., Hattersley, A. T., Jones, A. G., & MASTERMIND consortium (2018). Precision Medicine in Type 2 Diabetes: Clinical Markers of Insulin Resistance Are Associated With Altered Short- and Long-Term Glycemic Response to DPP-4 Inhibitor Therapy. *Diabetes Care*, 41(4), 705-712. <https://doi.org/10.2337/dc17-1827>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Supplementary material to:

Precision medicine in Type 2 diabetes: Clinical markers of insulin resistance are associated with altered short and long-term glycemic response to DPP4-inhibitor therapy

Authors: John M Dennis, Beverley M Shields, Anita V Hill, Bridget A Knight, Timothy J McDonald, Lauren Rogers, Michael N Weedon, William E Henley, Naveed Sattar, Rury R Holman, Ewan R. Pearson, Andrew T. Hattersley, Angus G. Jones on behalf of the MASTERMIND consortium

PRIBA Study Group

Lead Centre:

Royal Devon and Exeter NHS Foundation Trust/University of Exeter: Anita Hill, Rob Bolt, Jane Stewart, Bridget Knight, Tim McDonald, Beverley Shields, Angus Jones, Andrew Hattersley, Gayle Githens-Mazer, Tina Sanders, Kirsty Wensley

NIHR Clinical Research Network:

Ipswich Hospital NHS Trust: Gerry Rayman, Sue Hood, Jo Rosier, Jane Jiao, Debbie Simmonds, Caroline Calver

North Bristol Hospitals NHS Trust: Andrew Johnson, Sharon Tovey, Jade Bennet, Dafydd Wilson Evans, Philippa Lamb, Hilary Holloway, B Moore

Northampton General Hospital NHS Trust: Charles Fox, Kathy Hall, L James, C Smith

Northern Devon Healthcare NHS Trust: Alastair Watt, Geraldine Belcher, Amanda Skinner

Oxford Centre for Diabetes, Endocrinology and Metabolism: Steve Gough, Judy MacDonald, Lynne Nairn, Sue Rous

Plymouth Hospitals NHS Trust: Ann Millward, Margaret Blackmore, Migaila Watt

Portsmouth Hospitals NHS Trust: Mike Cummings, Sharon Allard, Elaine Hallett, Jane Rowney

Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust: David Kerr, Patricia Sanders, Carina Vickers

Royal Cornwall Hospitals NHS Foundation Trust: Steve Creely, Duncan Browne, Helen Chenoweth, Terri Chant, Sue Durkin

Royal Stoke University Hospital North Midlands: Ellen Hodgson, Gemma Reddell, Loretta Barnett, Jane Deleaney

South Devon Healthcare NHS Foundation Trust: Richard Paisey, Sue Bunce, Dawn Tomlinson, Mary Costello

South Warwickshire NHS Foundation Trust: Peter Horrocks, Penny Parsons, Alex Smith

Surrey and Sussex Healthcare NHS Trust: James Clark, Tracey Shewan, Louise Nimako


Taunton and Somerset NHS Foundation Trust: Rob Andrews, Catherine Thompson, Donna Archer

West Hertfordshire Hospitals NHS Trust: Thomas Galliford, Elaine Walker, Lynn Curry, Sindi Masuka, Cathy Constantin


Yeovil District Hospital NHS Foundation Trust: Seshadri Pramodh, Linda Balian, James Gibbons, Claire Buckley

Supplementary figure 1a: PRIBA study Profile

*Analysis inclusion criteria: no additional of non DPP4 glucose lowering therapy, or discontinuation of >1 co-therapy, between baseline and follow up HbA1c. Self-reported DPP4 adherence (over 2 weeks prior to HbA1c) $\geq 75\%$. Participant not receiving insulin treatment at study baseline.



Supplementary Figure 1b: CPRD study profile



*No change in co-therapy over period of interest

**Additional inclusion criteria: baseline HbA1c 7-11%, remained on DPP4i therapy for at least 3m

Supplementary table 1: Associations between markers of insulin resistance and HbA1c response after 6 months, overall (Data table for Figure 1) and stratified by sex in a) PRIBA and b) CPRD. Beta coefficient represents the change in HbA1c per standard deviation higher predictor level, a positive coefficient represents an association with reduced response.

a) PRIBA

| | All participants | | | Males (64%) | | Females (36%) | |
|-----------------------|---------------------------------|---------------------------|---------|---------------------------|---------|---------------------------|---------|
| | Number with valid baseline data | Beta coefficient (95% CI) | p-value | Beta coefficient (95% CI) | p-value | Beta coefficient (95% CI) | p-value |
| HOMA-IR* | 242 | 2.17 (0.62 to 3.72) | 0.01 | 1.69 (-0.26 to 3.64) | 0.09 | 2.87 (0.22 to 5.52) | 0.03 |
| Fasting C-peptide | 251 | 1.67 (0.17 to 3.17) | 0.03 | 1.44 (-0.49 to 3.36) | 0.14 | 1.91 (-0.56 to 4.39) | 0.13 |
| UCPCR** | 203 | 1.65 (-0.07 to 3.37) | 0.06 | 3.35 (1.05 to 5.65) | <0.01 | -0.56 (-3.16 to 2.04) | 0.67 |
| Fasting Triglycerides | 240 | 2.54 (0.99 to 4.08) | <0.01 | 2.20 (0.35 to 4.05) | 0.02 | 3.31 (0.33 to 6.29) | 0.03 |
| BMI | 254 | 0.96 (-0.54 to 2.46) | 0.21 | 1.23 (-0.89 to 3.35) | 0.25 | 0.29 (-1.98 to 2.56) | 0.80 |
| HDL | 243 | 0.20 (-1.36 to 1.75) | 0.81 | 0.56 (-1.1 to 2.62) | 0.60 | -0.95 (-3.60 to 1.70) | 0.48 |
| SHBG*** | 214 | -1.19 (-2.81 to 0.42) | 0.15 | -1.19 (-3.35 to 0.97) | 0.28 | -1.23 (-3.79 to 1.33) | 0.34 |

* HOMA2 measured insulin resistance **UCPCR = post meal urine C-peptide Creatinine ratio; ***SHBG = sex-hormone binding globulin

b) CPRD

| | All patients | | | Males (61%) | | Females (39%) | |
|---------------|---------------------------------|---------------------------|---------|---------------------------|---------|---------------------------|---------|
| | Number with valid baseline data | Beta coefficient (95% CI) | p-value | Beta coefficient (95% CI) | p-value | Beta coefficient (95% CI) | p-value |
| BMI | 19,430 | 0.96 (0.78 to 1.15) | <0.01 | 1.04 (0.78 to 1.30) | <0.01 | 0.78 (0.51 to 1.05) | <0.01 |
| Triglycerides | 15,404 | 0.72 (0.50 to 0.93) | <0.01 | 0.82 (0.56 to 1.09) | <0.01 | 0.57 (0.21 to 0.95) | <0.01 |
| HDL | 17,058 | -0.08 (-0.28 to 0.13) | 0.46 | -0.22 (-0.49 to 0.05) | 0.11 | -0.21 (-0.55 to 0.13) | 0.22 |

Supplementary table 2: Effect sizes for insulin resistance markers controlling for routine clinical characteristics (baseline HbA1c, age at therapy, sex, duration of diabetes, eGFR, ethnicity (CPRD only: white, non-white, missing) and co-therapy change (PRIBA only, CPRD patients all on unchanged therapy). Beta coefficient represents the change in HbA1c at 6 months per standard deviation increase in the predictor, a positive coefficient represents an association with reduced response.

a) PRIBA

| | Number of patients | Beta coefficient (95% CI) | p-value |
|-----------------------|---------------------------|----------------------------------|----------------|
| HOMA-IR | 205 | 2.57 (0.73 to 4.40) | <0.01 |
| Fasting C-peptide | 212 | 2.13 (0.33 to 3.94) | 0.02 |
| Fasting Triglycerides | 215 | 2.34 (0.44 to 4.25) | 0.02 |
| BMI | 215 | 0.67 (-1.15 to 2.50) | 0.47 |
| HDL-c | 215 | 0.09 (-1.69 to 1.86) | 0.92 |
| SHBG | 185 | -1.04 (-3.00 to 0.91) | 0.29 |

b) CPRD

| | Number of patients | Beta coefficient (95% CI) | p-value |
|---------------|---------------------------|----------------------------------|----------------|
| Triglycerides | 13,089 | 0.67 (0.41 to 0.94) | <0.01 |
| BMI | 11,683 | 0.87 (0.61 to 1.12) | <0.01 |
| HDL-c | 13,187 | -0.11 (-0.35 to 0.14) | 0.40 |

Supplementary table 3: Associations between triglycerides and BMI and HbA1c response after 6 months adjusted for baseline HbA1c, fasting glucose and co-therapy change in PRIBA. Beta coefficient represents the change in HbA1c at 6 months per standard deviation increase in the predictor, a positive coefficient represents an association with reduced response.

| | Number of patients | Beta coefficient (95% CI) | p-value |
|-----------------------|---------------------------|----------------------------------|----------------|
| HOMA-IR* | 242 | 1.76 (0.15 to 3.38) | 0.03 |
| Fasting C-peptide | 242 | 1.71 (0.21 to 3.21) | 0.03 |
| UCPCR** | 195 | 1.50 (-0.22 to 3.21) | 0.09 |
| Fasting Triglycerides | 231 | 2.39 (0.84 to 3.94) | <0.01 |
| BMI | 244 | 1.10 (-0.41 to 2.61) | 0.15 |
| HDL-c | 233 | 0.20 (-1.36 to 1.76) | 0.80 |
| SHBG*** | 206 | -0.90 (-2.53 to 0.73) | 0.28 |

* HOMA2 measured insulin resistance **UCPCR = post meal urine C-peptide Creatinine ratio;
 ***SHBG = sex-hormone binding globulin

Supplementary table 4: Associations between routine markers of insulin resistance and HbA1c response after 12 months in CPRD. For each predictor we ran a separate linear regression model, adjusted for baseline HbA1c. Beta coefficient represents the change in HbA1c at 12 months per standard deviation increase in the predictor, a positive coefficient represents an association with reduced response.

| | Number of patients | Beta coefficient (95% CI) | p-value |
|---------------|---------------------------|----------------------------------|----------------|
| Triglycerides | 13,942 | 0.70 (0.45 to 0.96) | <0.01 |
| BMI | 11,206 | 0.81 (0.59 to 1.04) | <0.01 |
| HDL-c | 12,273 | -0.32 (-0.57 to -0.08) | 0.01 |

Supplementary table 5: Associations between triglycerides and BMI and HbA1c response after 6 months in a combined model adjusted for baseline HbA1c, and (PRIBA only) co-therapy change in a) PRIBA and b) CPRD. Beta coefficient represents the change in HbA1c at 6 months per standard deviation increase in the predictor, a positive coefficient represents an association with reduced response.


| | a) PRIBA (n=240) | | b) CPRD (n=13,543) | |
|---------------|----------------------------------|----------------|----------------------------------|----------------|
| | Beta coefficient (95% CI) | p-value | Beta coefficient (95% CI) | p-value |
| Triglycerides | 2.33 (0.71 to 3.94) | <0.01 | 0.56 (0.32 to 0.79) | <0.01 |
| BMI | 0.44 (-1.17 to 2.04) | 0.59 | 0.96 (0.73 to 1.18) | <0.01 |

Supplementary table 6: CPRD hazard ratios for time to glycaemic failure (confirmed HbA1c ≥ 69 mmol/mol (8.5%)) for each predictor in the multivariable survival model (n=15,616)

| | Hazard ratio | 95% CI | p-value |
|--|---------------|-------------|---------|
| BMI & Triglyceride subgroup | | | |
| BMI<30 & TRGS<2.3 | 1 (reference) | | |
| BMI ≥ 30 OR TRGS ≥ 2.3 | 1.17 | 1.08-1.27 | <0.001 |
| BMI ≥ 30 & TRGS ≥ 2.3 | 1.28 | 1.16-1.41 | <0.001 |
| Clinical characteristics | | | |
| Baseline HbA1c (mmol/mol)* | 1.061 | 1.058-1.064 | <0.001 |
| Age at therapy start (year)* | 0.990 | 0.986-0.994 | <0.001 |
| Duration of diabetes (year)* | 1.004 | 0.997-1.011 | 0.29 |
| Female vs male sex | 1.075 | 0.999-1.152 | 0.04 |
| eGFR (ml/min/1.3m ²)* | 1.001 | 0.999-1.002 | 0.49 |

*For continuous variables the hazard ratio represents the change in hazard ratio for a 1 unit increase in the predictor. A hazard ratio > 1 indicates a higher value of that variable is associated with shorter durability of glycaemic response

Supplementary Figure 2: Probability of glycemic failure (confirmed HbA1c $\geq 8.5\%$) over 3 years in CPRD in subgroups defined by the presence or absence of obesity (BMI ≥ 30 kg/m²) and high triglycerides (TRGs ≥ 2.3 mmol/L) - Subgroup A: non-obese and normal triglycerides, Subgroup B: non-obese OR normal triglycerides, Subgroup C: obese and high triglycerides).



Supplementary table 7: CPRD hazard ratios by BMI & Triglyceride subgroup for time to glycaemic failure defined as

a) Confirmed HbA1c ≥ 53 (7.5%) (n=15,616)*

| | Hazard ratio | 95% CI | p-value |
|--|---------------------|---------------|----------------|
| BMI & Triglyceride subgroup | | | |
| BMI<30 & TRGS<2.3 | 1 (reference) | | |
| BMI \geq 30 OR TRGS \geq 2.3 | 1.08 | 1.02-1.14 | 0.01 |
| BMI \geq 30 & TRGS \geq 2.3 | 1.17 | 1.09-1.25 | <0.001 |

b) Confirmed return to baseline HbA1c level specific to each patient (n=15,616)*

| | Hazard ratio | 95% CI | p-value |
|--|---------------------|---------------|----------------|
| BMI & Triglyceride subgroup | | | |
| BMI<30 & TRGS<2.3 | 1 (reference) | | |
| BMI \geq 30 OR TRGS \geq 2.3 | 1.14 | 1.05-1.24 | 0.002 |
| BMI \geq 30 & TRGS \geq 2.3 | 1.29 | 1.16-1.42 | <0.001 |

*adjusted for baseline HbA1c, age at therapy, duration of diabetes, sex and eGFR

GLP-1 receptor agonist comparison analysis

Supplementary table 8: GLP-1 receptor agonists - subject baseline characteristics

| | | PRIBA (n=339) | CPRD (n=4,464) |
|---|--------------|------------------------------|---------------------------|
| Characteristics | | | |
| mean (SD) unless stated | | | |
| Baseline HbA1c (mmol/mol) | | 83 (18) | 79 (17) |
| Baseline HbA1c (%) | | 9.7 (1.7) | 9.4 (1.6) |
| Age at therapy start (years) | | 55 (10) | 59 (9) |
| Age at diagnosis (years) | | 47 (10) | 51 (8) |
| Male sex (%) | | 56% | 58% |
| Duration of diabetes (years) | | 8 (5) | 8 (5) |
| BMI - median (IQR); mean(SD) | | 38 (35-44); 40 (8) | 37 (34-42); 38 (7) |
| Ethnicity (%) | | | |
| | White | 95% | 46% |
| | Non-White | 5% | 3% |
| | Missing | 0% | 51% |
| Biomarkers | | | |
| median (IQR); mean (SD) unless stated | | | |
| *=log-transformed | | | |
| Triglycerides (mmol/L) | | 1.9 (1.4-2.6); 2 (1.1)* | 2.0 (1.4-2.8); 2.0 (0.5)* |
| HDL-c (mmol/L) | | 1.1 (0.9-1.2); 1.1 (0.3)* | 1.1 (0.9-1.2); 1.1 (0.2)* |
| LDL-c (mmol/L) | | 2.2 (1.8-2.8); 2.2 (0.8)* | 2.0 (1.6-2.6); 2.0 (0.4)* |
| SHBG (nmol/L) | | 23 (17-36); 24 (14)* | NA |
| Fasting C-peptide (pmol/L) | | 1310 (962-1700); 1255 (594)* | NA |
| HOMA2-%B | | 49 (32-75); 48 (29)* | NA |
| HOMA2 IR | | 4.1 (3.0-5.3); 4.0 (2.1)* | NA |
| UCPCR nmol/mmol | | 3.5 (1.9-6.0); 3.1 (2.8)* | NA |
| eGFR (ml/min/1.3m ²) | | 92 (77-111); 95 (27) | 88 (74-102); 88 (22) |
| GAD or IA2 positive (%) | | 1% | NA |
| Therapy | | | |
| Number of concomitant therapies at therapy start (% of total) | | | |
| | 0 | 1% | 2% |
| | 1 | 26% | 36% |
| | 2 | 51% | 54% |
| | 3+ | 22% | 8% |
| GLP-1 type (% of total) | | | |
| | Dulaglutide | 0% | 1% |
| | Exenatide | 40% | 44% |
| | Liraglutide | 60% | 50% |
| | Lixisenatide | 0% | 6% |


Supplementary table 9: GLP-1 receptor agonists - Associations between markers of insulin resistance and HbA1c response after 6 months, overall (Data table for Supplementary Figure 2) in a) PRIBA and b) CPRD. Beta coefficients represent the change in HbA1c per standard deviation higher predictor level, a positive coefficient represents an association with reduced response.

| | PRIBA | | | CPRD | | |
|-------------------|---------------------------------------|------------------------------|---------|---------------------------------------|------------------------------|---------|
| | Number with valid baseline data | Beta coefficient (95% CI) | p-value | Number with valid baseline data | Beta coefficient (95% CI) | p-value |
| HOMA-IR* | 300 | 0.73 (-0.99 to 2.44) | 0.41 | NA | NA | NA |
| Fasting C-peptide | 333 | 0.24 (-1.38 to 1.85) | 0.78 | NA | NA | NA |
| UCPCR** | 245 | 0.45 (-1.37 to 2.26) | 0.63 | NA | NA | NA |
| Triglycerides | 316 | 0.07 (-1.59 to 1.73) | 0.93 | 2,848 | -0.07 (-0.65 to 0.49) | 0.80 |
| BMI | 337 | 0.11 (-1.49 to 1.70) | 0.90 | 4,016 | -0.19 (-0.66 to 0.28) | 0.43 |
| HDL | 315 | 1.08 (-0.55 to 2.71) | 0.19 | 3,126 | 0.01 (-0.53 to 0.55) | 0.97 |
| SHBG*** | 317 | 0.56 (-1.09 to 2.21) | 0.50 | NA | NA | NA |

Supplementary table 10: Mean (95% CI) 6 month baseline HbA_{1c} standardised glycemic response (mmol/mol) for DPP4-inhibitors and GLP receptor agonists in PRIBA and CPRD. Baseline HbA_{1c} is standardised to the mean PRIBA baseline level of 74mmol/mol (8.9%) for all subgroups.

| BMI & Triglyceride subgroup | DPP4-inhibitors | | GLP-1 receptor agonists | |
|-----------------------------|-------------------------|--------------------------|---------------------------|-------------------------|
| | PRIBA | CPRD | PRIBA | CPRD |
| BMI<30 & TRGS<2.3 | -11.26 (-14.1 to -8.43) | -10.28 (-10.72 to -9.85) | -9.64 (-22.75 to 3.48) | -9.49 (-12.2 to -6.78) |
| BMI≥30 OR TRGS≥2.3 | -9.94 (-12.24 to -7.64) | -8.79 (-9.12 to -8.47) | -14.7 (-17.11 to -12.29) | -10.16 (-10.91 to -9.4) |
| BMI≥30 & TRGS≥2.3 | -5.23 (-8.62 to -1.84) | -7.47 (-7.95 to -6.99) | -13.54 (-16.69 to -10.39) | -9.64 (-10.65 to -8.63) |

Supplementary Figure 3: GLP-1 receptor agonists - predicted mean absolute HbA_{1c} change from baseline at 6 months in a) PRIBA b) CPRD across subgroups defined by the presence or absence of obesity (BMI≥30 kg/m²) and high triglycerides (TRGs ≥2.3mmol/L) - *Subgroup A*: non-obese and normal triglycerides, *Subgroup B*: non-obese OR normal triglycerides, *Subgroup C*: obese and high triglycerides. Baseline HbA_{1c} is standardised to the mean PRIBA baseline level of 74mmol/mol (8.9%) for all subgroups. Error bars denote 95% confidence intervals.



Supplementary table 10: GLP-1 receptor agonists - CPRD hazard ratios for time to glycaemic failure (confirmed HbA1c ≥ 69 mmol/mol (8.5%)) for each predictor in the multivariable survival model (n=2,795)

| | Hazard ratio | 95% CI | p-value |
|-----------------------------------|---------------------|---------------|----------------|
| BMI* | 1.01 | 1.00-1.02 | 0.29 |
| Triglycerides* | 0.99 | 0.95-1.04 | 0.80 |
| Baseline HbA1c (mmol/mol)* | 1.05 | 1.04-1.06 | <0.001 |
| Age at therapy start (year)* | 0.98 | 0.97-0.99 | <0.001 |
| Duration of diabetes (year)* | 1.01 | 0.99-1.03 | 0.21 |
| Female vs male sex | 0.78 | 0.67-0.91 | 0.001 |
| eGFR (ml/min/1.3m ²)* | 1.00 | 1.00-1.00 | 0.63 |

*For continuous variables the hazard ratio represents the change in hazard ratio for a 1 unit increase in the predictor. A hazard ratio > 1 indicates a higher value of that variable is associated with shorter durability of glycaemic response

Supplementary table 11: GLP-1 receptor agonists - CPRD hazard ratios for time to glycaemic failure (confirmed HbA1c ≥ 69 mmol/mol (8.5%)) for each predictor in the multivariable survival model (n=2,795)

| | Hazard ratio | 95% CI | p-value |
|--|---------------|-----------|---------|
| BMI & Triglyceride subgroup | | | |
| BMI<30 & TRGS<2.3 | 1 (reference) | | |
| BMI ≥ 30 OR TRGS ≥ 2.3 | 1.21 | 0.85-1.73 | 0.29 |
| BMI ≥ 30 & TRGS ≥ 2.3 | 1.13 | 0.79-1.63 | 0.50 |
| Clinical characteristics | | | |
| Baseline HbA1c (mmol/mol)* | 1.05 | 1.04-1.06 | <0.001 |
| Age at therapy start (year)* | 0.98 | 0.97-0.99 | <0.001 |
| Duration of diabetes (year)* | 1.01 | 0.99-1.03 | 0.22 |
| Female vs male sex | 0.79 | 0.68-0.92 | 0.002 |
| eGFR (ml/min/1.3m ²)* | 1.00 | 1.00-1.00 | 0.60 |

*For continuous variables the hazard ratio represents the change in hazard ratio for a 1 unit increase in the predictor. A hazard ratio > 1 indicates a higher value of that variable is associated with shorter durability of glycaemic response

Supplementary Figure 4: GLP-1 receptor agonists - Probability of glycaemic failure (confirmed HbA1c $\geq 8.5\%$) over 3 years in CPRD in subgroups defined by the presence or absence of obesity (BMI ≥ 30 kg/m²) and high triglycerides (TRGs ≥ 2.3 mmol/L) - Subgroup A: non-obese and normal triglycerides, Subgroup B: non-obese OR normal triglycerides, Subgroup C: obese and high triglycerides).

